

(12) **UK Patent Application** (19) **GB** (11) **2 248 618** (13) **A**
 (43) Date of A publication **15.04.1992**

(21) Application No **9124015.0**

(22) Date of filing **15.06.1989**

Date lodged **12.11.1991**

(30) Priority data

(31) **8814458**

(32) **17.06.1988**

(33) **GB**

(62) Derived from Application No **8913815.0** under Section 15(4) of the Patents Act 1977

(71) Applicant

John Wyeth & Brother Limited

(Incorporated in the United Kingdom)

**Huntercombe Lane South, Taplow, Maidenhead,
Berkshire, SL6 0PH, United Kingdom**

(72) Inventor

Robin Gerald Shepherd

(74) Agent and/or Address for Service

Dr D F Wileman

**Wyeth Laboratories, Huntercombe Lane South, Taplow,
Maidenhead, Berks, SL6 0PH, United Kingdom**

(51) INT CL⁵

C07D 215/18 221/04

(52) UK CL (Edition K)

**C2C CAB CZH C1534 C1544 C213 C247 C25Y
C250 C251 C305 C31Y C314 C337 C440 C776
U1S S1347**

(56) Documents cited

GB 2219798 A

(58) Field of search


UK CL (Edition K) **C2C CBE CWE CZH**

INT CL⁵ **C07D 215/18 221/04**

Online databases: **CAS-ONLINE**


(54) **Process for preparing bromo-pyridine derivatives**

(57) Compounds of formula



(IVa)

wherein R¹ and R² each independently represent hydrogen, lower alkyl, lower alkoxy, carboxyloweralkyl, carboxy, hydroxyloweralkyl, halogen, haloloweralkyl, lower alkoxycarbonyl, optionally substituted aryl or optionally substituted aralkyl, n represents an integer from 3 to 6; R³ represents hydrogen or single or multiple substitution on one or more of the aliphatic carbons by one or more substituents selected from lower alkyl, optionally substituted aryl and optionally substituted aralkyl, are prepared by reacting a compound of formula



(VII)

with either HBr and bromine or with bromine in an inert solvent.


The date of filing shown above is that provisionally accorded to the application in accordance with the provisions of Section 15(4) of the Patents Act 1977 and is subject to ratification or amendment.

GB 2 248 618 A

HETEROCYCLIC COMPOUNDS


This invention relates to heterocyclic compounds, more particularly to pyridine derivatives, processes for preparing them, pharmaceutical compositions containing them and to intermediates therefor.

5 This invention provides compounds of formula



or a salt thereof, wherein R^1 and R^2 each independently represent hydrogen, alkyl, alkoxy, carboxyalkyl, carboxy, hydroxyalkyl, halogen, haloalkyl, alkoxycarbonyl, optionally substituted aryl or
 10 optionally substituted aralkyl, n represents an integer from 3 to 6; R^3 represents hydrogen or single or multiple substitution on one or more of the aliphatic carbons, eg. by one or more substituents selected from lower alkyl, optionally substituted aryl and optionally
 15 substituted aralkyl;

A represents a group of formula (i) or (ii) below:



in which R^4 , R^6 and R^7 each independently represent hydrogen or lower alkyl (providing that when R^5 is NH_2 R^4 is hydrogen); m is 0 or 1;
 20 R^5 represents hydrogen, NH_2 , OH or loweralkoxy, and X is =O, =NH or =NOH, the attachment of A to B being from either end;

and B represents an optionally substituted aryl or heteroaryl radical.

By the term aryl as used herein is meant any monovalent carbocyclic radical possessing aromatic character and includes groups having 6 to 10 carbon atoms such as phenyl and naphthyl. By the term heteroaryl as used herein is meant any monovalent heterocyclic group possessing aromatic character and includes groups having 5 to 10 ring atoms and one or more heteroatoms selected from oxygen, nitrogen and sulphur. Examples of heteroaryl radicals are furyl, thienyl, pyrrolyl, pyridinyl, pyrimidinyl, quinolyl, isoquinolyl, benzimidazolyl, thiazolyl and imidazolyl.

The term alkyl when used to signify a group or part of a group such as hydroxyalkyl or aralkyl means any straight or branched saturated aliphatic hydrocarbon especially those having 1 to 6 carbon atoms, eg. -C_4 carbon atoms. Examples are methyl, ethyl, n-propyl, isopropyl, n-butyl, n-pentyl and n-hexyl.


By the term 'optionally substituted' is meant optional substitution on carbon atoms by one or more substituents, eg. substituents commonly used in pharmaceutical chemistry, eg. halogen (eg. Cl, Br, F), alkyl, alkyloxy, haloalkyl (eg. CF_3), or haloalkoxy (eg. $\text{CHF}_2\text{O-}$, $\text{CF}_3\text{CH}_2\text{O-}$), NO_2 , NH_2 , CN, alkylamino, dialkylamino, carboxy, alkyloxycarbonyl, acyl, acylamino, aryl, (eg. phenyl) or aminoalkyl.

The term 'lower' as used herein to qualify a group means such a group contains 1 to 6 carbon atoms.


-4-

Examples of the groups R^1 and R^2 are independently hydrogen, lower alkyl (eg. methyl, ethyl, n-propyl, isopropyl, n-butyl, n-pentyl and n-hexyl), $-COOH$, $-COO$ lower alkyl (eg. $COOMe$, $COOEt$), $-CH_2OH$, Br, Cl, CF_3 , OMe, optionally substituted phenyl or optionally substituted benzyl.


Preferred values of n are 3, 4, or 5 such that the compounds have one of the following formulae



(Ia)



(Ib)



(Ic)

Compounds of formula Ic are most preferred.

Examples of R^3 are one or more substituents selected from methyl, ethyl, methoxy, ethoxy, phenyl, phenyl substituted by halogen, lower alkyl or lower alkoxy, benzyl and benzyl substituted by halogen, lower alkyl or lower alkoxy. Examples of multisubstitution on one carbon atom is gem-dimethyl.

Examples of R^4 , R^6 and R^7 are independently hydrogen, methyl and ethyl.


The linkage of $-A-$ to the pyridine ring and B can be either way round so that examples of $-A-B$ are $-CHOH-B$,

-5-

-CHOHCH₂-B, -CH₂CHOH-B, -CO-B, -COCH₂B, -CH₂COB,
 -CH(OCH₃)-B, -CH(OCH₃)CH₂B and -CH₂CH(OCH₃)-B.
 Preferably A is -CHOH- or -CO-.

Examples of B are phenyl, pyridyl (eg. pyrid-2-yl),
 5 1-naphthyl, 2-naphthyl which groups may be substituted
 as defined herein, for example by one or more
 substituents selected from lower alkyl, lower alkoxy,
 halogen, haloloweralkyl, nitro, amino, lower
 alkylamino, cyano, carboxy, lower alkoxy carbonyl and
 10 lower alkyl carbonyl.

In a preferred aspect this invention provides compounds
 of formula



(Id)

or salts thereof,

wherein n represents 4 or 5; R⁸ represents lower alkyl,
 15 halogen, carboxy, loweralkoxy; R⁹ represents hydrogen
 or a lower alkyl group; A¹ represents -CO- or -CHOH-
 and B¹ represents optionally substituted phenyl,
 especially where the substituent is selected from one
 or more of the following: lower alkyl, (eg. o-,m-or p-
 20 methyl); halogen (eg. o-,m-or p-chloro or bromo); lower
 alkoxy (eg. o-,m- or p-methoxy), and carboxy, eg. o-,m-
 or p- carboxy. Other examples of B¹ (and B) include
 2,4-dimethylphenyl; 3,4-dimethylphenyl; 4-ethylphenyl;

-6-

4-isopropylphenyl, 4-isobutylphenyl,
2-methoxy-4-methylphenyl, 2-chloro-4-methylphenyl.

The compounds of formula I can possess one or more
asymmetric centres and hence optical isomers and
5 mixtures thereof are possible, eg. when A is CHOH or
R³ represents monosubstitution. All such isomers and
mixtures thereof are included within the scope of this
invention. Where any reaction process produces
mixtures of such isomers standard resolution techniques
'0 may be applied to separate a specific isomer.

The compounds of formula I may be obtained in free base
form or as acid addition salts as desired. Examples of
such salts include salts with pharmaceutically
acceptable acids such as hydrochloric, hydrobromic,
'5 hydroiodic, sulphuric, phosphoric, nitric, acetic,
citric, tartaric, fumaric, succinic, malonic, formic,
maleic acid or organosulphonic acids such as methane
sulphonic or p-toluene sulphonic acids.

When acidic substituents are present it is also
20 possible to form salts with bases eg. alkali metals
(such as sodium) or ammonium salts. Such salts of the
compounds of formula I are included within the scope of
this invention.

When basic substituents are present then quaternary
25 ammonium salts may be formed by quaternizing with an
alkylating agent such as alkyl, or alkyl halides.

The compounds of formula I possess pharmaceutical
activity in standard tests, in particular
anti-inflammatory activity. Anti-inflammatory

-7-

compounds of the non-steroidal type currently available generally have a tendency to cause gastric irritation and some are markedly ulcerogenic. It has surprisingly been found that in standard tests compounds of formula
5 I appear to be devoid of ulcerogenic properties even at high doses.

The compounds of formula I were tested for anti-inflammatory activity in the following test procedures:

Procedure 1

10 Anti-inflammatory activity of a compound is assessed by its ability to inhibit experimentally induced edema in the hind paw of the rat.

Groups of six male Sprague-Dawley rats (Charles River), weighing between 150-165 g, were used in these
15 experiments. Drugs were administered p.o. in 0.5% methylcellulose (400 centipoise). One hour after administering drugs or vehicle, 0.1 ml of 1% carrageenan was injected subplantar into the right hind paw. Right hind paw volumes (ml) were measured prior
20 to carrageenan injection using a mercury plethysmograph (ie. zero time reading). After three hours, the right hind paw volumes were re-measured and paw edema was calculated for each rat by subtracting the zero time reading from the three hour reading. The percent
25 change in paw edema was calculated according to the formula:

$$\% \text{ Inhibition} = 100\% \times$$

$$\frac{\text{Mean vol. swelling control} - \text{mean vol. swelling of test}}{\text{Mean vol. swelling of control}}$$

The Dunnett's test was used to determine statistical significance ($P < 0.05$).

In the above test representative compounds of formula I gave the results shown in TABLE I.

TABLE 1

	Compounds of <u>formula I</u> <u>Dose 50mgs/kg</u>	Mean Paw Edema @ 3 <u>hours ml +S.E</u>	Number of <u>Assays</u>	Average % change @ <u>+ 3 hours</u>
5	Control	1.1 \pm 0.05	100	-
		0.45 \pm 0.06	2	-59
	1-(6,7,8,9-Tetra hydro-3-methyl-5H- cyclohepta[b]- pyrid-2-yl)-1- 10 (4-methylphenyl) methanol (Example 1)			
	6,7,8,9,-Tetrahydro -3-methyl-2-(methyl- benzoyl)-5H-cyclo- 15 hepta[b]pyridine, hydrochloride (Example 2)	0.47 \pm 0.05	2	-58
	1-(6,7,8,9-Tetra- hydro-5H-cyclohepta 20 [b]pyrid-2-yl)- -1-(4-methylphenyl) methanol (Example 3)	0.83 \pm 0.06	2	-24
	1-(6,7,8,9-Tetrahydro- 3-methyl- 25 -5H-cyclohepta- [b]pyrid-2- -yl)-1-phenylmethanol	0.51 \pm 0.08	2	-54

-9-

Compounds which inhibit the swelling by about 20% are considered of interest. The results show that all the compounds were active with the compounds of Examples 1,3 and 4 demonstrating marked anti-inflammatory activity at the dose level tested.

Procedure 2

In this procedure polyarthrititis is induced in male Lewis strain rats by the injection of a tubercle bacilli in the subplanter tissue of the right hind paw.

Groups of ten male Lewis rats (Charles River), weighing between 150-170 g, were injected s.c. into the right hind paw with dessicated Mycobacterium butyricum (0.5 mg/0.1 ml) suspended in light mineral oil. Drugs were administered orally in 0.5% methylcellulose from day 0 to 15 (except weekends). Both hind paw volumes (ml) were measured by mercury plethysmography at the time of injection of adjuvant (day 0). Paw volumes were measured on day 4 (injected paw only) and on day 16 (uninjected paw) to determine the nonspecific and immunologically-induced inflammation, respectively. Drug effects were expressed as a percentage change from vehicle-treated arthritic controls.

Active compounds will either prevent or reverse the joint swelling and associated sequella of polyarthrititis. A % change of about 20% from control is regarded as noteworthy.

In this test in a series of experiments compounds of formula I gave the results shown in Table 2.

-10-

TABLE 2

	<u>Compound of formula I</u>	<u>Dose level</u>	<u>% Change from Control</u>
	1-(6,7,8,9-tetrahydro-3-methyl-5H-cyclohepta[b]-pyrid-2-yl)-1-(4-methylphenyl)methanol	30 mg/kg	-47%
5	6,7,8,9-tetrahydro-3-methyl-2-(4-methylbenzoyl)-5H-cyclohepta[b]pyridine, hydrochloride	30 mg/kg	-26%
10	1-(6,7,8,9-tetrahydro-5H-cyclohepta[b]-pyrid-2-yl)-1-(4-methylphenyl)methanol	30 mg/kg	-33%
15	1-(6,7,8,9-tetrahydro-3-methyl-5H-cyclohepta[b]-pyrid-2-yl)-1-phenylmethanol	30 mg/kg	-45%

The results in Table 2 show the above-mentioned compounds of formula I exhibit marked ant inflammatory activity in the test.


In standard tests investigating gastric irritation representative compounds of formula I, namely 1-(6,7,8,9-tetrahydro-3-methyl-5H-

-11-

cyclohepta[b]pyrid-2-yl)-1-(4-methylphenyl)methanol and 6,7,8,9-tetrahydro-3-methyl-2-(4-methylbenzoyl)-5H-cyclohepta[b]pyridine, hydrochloride did not show any effect at the highest dose level tested of 300 mg/kg.


- 5 The compounds of formula I may be prepared by various methods all of which are included in the scope of this invention.

A first method for preparing compounds of formula I wherein A is $-\text{CR}^4\text{R}^5-(\text{CR}^6\text{R}^7)_m$ -where R^5 is OH and R^4 , R^6 , R^7 and m are as defined above comprises reacting a
 10 compound of formula .



(II)

- wherein R^1 , R^2 , R^3 and n are as defined above, p is 0 or 1 and M is sodium, potassium, lithium, Mg⁺hal where hal is halogen, eg. chlorine, bromine or iodine
 15 providing that hal is not chlorine when p is 0, with a compound of formula




(III)

- wherein B, R^3 , R^6 and R^7 are as defined above and q is 0 or 1 providing that p + q is 0 or 1, followed by protonation.

- 20 The reaction is conveniently carried out under anhydrous conditions in an appropriate inert solvent

- eg. tetrahydrofuran, toluene, diethyl ether, an alkane (such as hexane) or a cycloalkane (eg. cyclohexane) or mixtures of such solvents at or below room temperature. Preferably the starting material of formula II is
- 5 prepared in situ and the same solvent used for reaction with the compound of formula III. The final step in the reaction is to protonate the intermediate complex, eg. using water alcohol or an acid. In the
- 10 aforementioned reaction any carbon atom epi to the nitrogen (for example the 9-position when n is 5) may be protected prior to reaction for example by using one or two organic silyl protecting groups, eg. trialkylsilyl such as trimethylsilyl and removing the protecting group(s) after the reaction.
- 15 The starting material of formula II may generally be prepared by metallation reactions from appropriate corresponding compounds of formula




- optionally protected in the epi position where Z is hydrogen or halogen and n, p, R¹, R², R³ and R⁶ and R⁷ are
- 20 as defined above. For example the starting material of formula II wherein M is lithium, sodium or potassium may be prepared by reacting a compound of formula IV wherein Z is hydrogen or bromine or iodine with a metal alkyl, eg. MR¹⁰ wherein M is Li, Na or K and R¹⁰ is
- 25 alkyl, aryl or aralkyl, for example n-butyllithium. Compounds of formula II where M is Mg^{hal} may be prepared by treating a compound of formula IV wherein Z is halogen with magnesium under conditions known for the

-13-

preparation of Grignard reagents.

Compounds of formula IV are known compounds or can be made by known methods for analogous compounds. For example 5,6,7,8-tetrahydroquinolines can be prepared by reacting 3-aminoacroleins with cyclohexanones according to the method of Breitmaier and Bayer (Tet Letts 1970, 38, 3291). Similarly 6,7,8,9-tetrahydro-5H-cyclohepta[b]pyridines and cyclopenta[b]pyridines can be prepared by reacting respective cycloheptanone or cyclopentanones with 3-aminoacroleins (eg. 3-amino-2-methylacrolein). These and other methods are described in UK Patent Specification 1,432,378.

Another aspect of this invention provides a new and particularly convenient route to compounds of formula IV wherein p is 0 and Z represents bromine. Such compounds may be prepared by reacting a compound of formula



(VII)


wherein n, R¹, R² and R³ are as defined above with either bromine or HBr and bromine in an inert solvent, such as an alkanolic acid eg. acetic acid or halocarbon, eg. methylene dichloride. This is believed to be the first direct cyclisation route to 2-bromo-cycloalkeno-pyridines. Previously such compounds were prepared by halogenating a corresponding 2-hydroxypyridine. The new process provides a particularly useful route via

-14-

the appropriate intermediates ultimately to compounds of formula I wherein $n=5$. The reaction may be carried out without heating if desired using approximately equimolar quantities of bromine and compound of formula VII. When HBr is used the amount may vary from a catalytic amount to a molar excess, eg. 5M.


A further aspect of this invention provides intermediates of formulae II and IV wherein n is 5

Compounds of formula I wherein A is $-\text{CX}-(\text{CR}^6\text{R}^7)_m-$ where X is =NH and m , R^6 and R^7 are as defined above may be prepared by reacting a compound of formula:



(V)

(if required protected in the epi position) wherein n , R^1 , R^2 and R^3 are as defined above, p is 0 or 1 and M^1 is lithium, sodium, potassium or Mg hal where hal is chlorine, bromine or iodine, with a nitrile of formula:



(VI)

wherein B, R^6 and R^7 are as defined above and q is 0 or 1 providing that $p + q = 0$ or 1 and if required removing any protecting group. The above reaction may be conveniently carried out at or below room temperature in an inert solvent under anhydrous conditions.

Compounds of formula I wherein A is $-\text{CR}^4\text{R}^5-(\text{CR}^6\text{R}^7)_m-$

-15-

wherein R^5 is amino, R^4 is hydrogen and m, R^6 and R^7 are as defined above, may be prepared by a process which comprises reducing a corresponding compound of formula I wherein A is $-CX-(CR^6R^7)_m$ - where X is = NH. The
5 reduction is conveniently carried out using an alkali metal borohydride, such as sodium borohydride in alcohol solvent, eg. ethanol.

Compounds of formula I wherein A is $-CO-(CR^6R^7)_m$ - where m, R^6 and R^7 are as defined above may be prepared
10 by a process which comprises
a) hydrolysing a compound of formula I wherein A is $-CX-(CR^6R^7)_m$ - where X is = NH or = NOH,
or (b) oxidising a compound of formula I wherein A is
15 $-CR^4R^5-(CR^6R^7)_m$ - wherein R^4 is hydrogen, R^5 is OH and m, R^6 and R^7 are as hereinbefore defined. The hydrolysis step (a) may be carried out using an aqueous acid with heating. The oxidation step (b) may be carried out using an oxidation agent known for
20 oxidising alcohols to ketones, eg. manganese dioxide.

Where the product of a reaction is a ketone then the oxime derivative thereof may be prepared in the usual manner eg. using hydroxylamine. Such derivatives may be converted to the ketone by hydrolysis.


25 When a compound of formula I is prepared in which R^5 is OH then that compound may be converted to a compound of formula I wherein R^5 is -O- lower alkyl by processes known for lower alkyl etherification (eg. Williamson Synthesis, Merck Index 10th Edition page
30 ONR-96 and references cited therein). For example the lower alkyl ether may be prepared by first forming the

-16-

alkali metal salt of the alcohol using an alkali metal hydride, then reacting with an alkyl halide, eg. methyl iodide.

Compounds of formula I wherein R^5 is hydrogen can be prepared by methods disclosed in UK Patent Specification 1,432,378 or by analogous methods. They may also be prepared by reducing corresponding alcohols using hydrogenation techniques. For example compounds of formula I wherein R^5 is OH may be reduced by hydrogenation using palladium charcoal to give the corresponding compound of formula I wherein R^5 is hydrogen.

Compounds of formula II which are useful as intermediates are novel compounds within the scope of this invention. Included within these intermediates are those having formula II in which the carbon epi to the nitrogen is protected by one or two organic silyl protecting groups eg. by trialkylsilyl. Accordingly the intermediates can be represented by formula IIa



wherein n, p, R^1, R^2, R^3, R^6 and R^7 are as hereinbefore defined, M is Na, K, Li or $MgHal$ where Hal is a halogen, and P^1 and P^2 each represent optional organic silyl protecting group on the carbon epi to the nitrogen.

-17-

Starting materials for the processes described herein are known compounds or can be prepared by analogous methods for known compounds.

5 This invention also provides pharmaceutical compositions comprising a compound of formula I or a pharmaceutically acceptable salt thereof.

For the pharmaceutical compositions any suitable carrier known in the art can be used. In such a composition, the carrier may be a solid, liquid or
10 mixture of a solid and a liquid. Solid form compositions include powders, tablets and capsules. A solid carrier can be one or more substances which may also act as flavouring agents, lubricants, solubilisers, suspending agents, binders, or tablet
15 disintegrating agents; it can also be encapsulating material. In powders the carrier is a finely divided solid which is in admixture with the finely divided active ingredient. In tablets the active ingredient is mixed with a carrier having the necessary binding
20 properties in suitable proportions and compacted in the shape and size desired. The powders and tablets preferably contain from 5 to 99, preferably 10-80% of the active ingredient. Suitable solid carriers are magnesium stearate, talc, sugar, lactose, pectin,
25 dextrin, starch, gelatin, tragacanth, methyl cellulose, sodium carboxymethyl cellulose, a low melting wax and cocoa butter. The term "composition" is intended to include the formulation of an active ingredient with encapsulating material as carrier, to give a capsule in
30 which the active ingredient (with or without other carrier) is surrounded by carriers, which is thus in association with it. Similarly cachets are included.

-18-

Sterile liquid form compositions include sterile solutions, suspensions, emulsions, syrups, and elixirs. The active ingredient can be dissolved or suspended in a pharmaceutically acceptable carrier, such a sterile
5 water, sterile organic solvent or a mixture of both. The active ingredients can be often be dissolved in a suitable organic solvent, for instance aqueous propylene glycol containing from 10 to 75% of the glycol by weight is generally suitable. Other
10 compositions can be made by dispersing the finely-divided active ingredient in aqueous starch or sodium carboxymethyl cellulose solution, or in a suitable oil, for instance arachis oil. Preferably the pharmaceutical composition is in unit
15 dosage form, the composition is sub-divided in unit doses containing appropriate quantities of the active ingredient; the unit dosage form can be a packaged composition, the package containing specific quantities of compositions, for example packeted powders or vials
20 or ampoules. The unit dosage form can be a capsule, cachet or tablet itself, or it can be the appropriate number of any of these in packaged form. The quantity of active ingredient in a unit dose of composition may be varied or adjusted from 10 to 500 mg or more, eg.
25 25 mg to 250 mg, according to the particular need and the activity of the active ingredient. The invention also includes the compounds in the absence of carrier wherethe compounds are in unit dosage form. Based on the results from animal studies the dosage range for
30 the treatment of humans using a compound of formula I will be in the range from about 5 mg to 2 g per day depending on the activity of the compound. The following Examples illustrate the invention and methods for preparing compounds of the invention.

-19-

EXAMPLE 11-(6,7,8,9-Tetrahydro-3-methyl-5H-cyclohepta[b]pyrid-2-yl)-1-(4-methylphenyl)methanol

a) A 1.36M solution of n-butyl lithium in hexane (19ml) in THF (20ml) was treated with a solution of 6,7,8,9-tetrahydro-3-methyl-5H-cyclohepta[b]pyridine (4.0g, 0.025m) in tetrahydrofuran (THF) (5ml) at 0°C. After 5 0.5 hours the solution was blown over, under nitrogen pressure, into a solution of trimethylsilyl chloride (10ml) in THF (30ml) cooled to 0°C. After 0.5 hours the solvent was removed under reduced pressure and the residue treated with hexane, filtered and evaporated to 10 give an oil containing 6,7,8,9-tetrahydro-3-methyl-9-trimethylsilyl-5H-cyclohepta[b]pyridine.

b) The oil from step (a) was dissolved in THF (50ml) and a 1.36M solution of n-butyl lithium in hexane (19.1) was added at 0°C. The mixture was stirred at 15 this temperature for a further 0.5 hours. 4-Methylbenzaldehyde (3.6g, 0.03m) in THF (5ml) was added rapidly at 0°C and the mixture stirred for a further 0.25 hours. A solution of 2M hydrochloric acid (60ml) was added and stirred for 0.25 hours, then 20 separated. The aqueous phase was separated and basified with solid potassium carbonate and then extracted with chloroform. The combined chloroform extracts were washed with water, dried (MgSO₄) and evaporated. The residue was treated with ether and the 25 crystallised material collected and dried to give the title compound (2.2g) m.p. 143-5°C.

Analysis:

Found C, 81.4; H, 8.3; N, 4.7.

C₁₉H₂₃NO requires C, 81.1; H, 8.2; N, 5.0%.

-20-

EXAMPLE 26,7,8,9-Tetrahydro-3-methyl-2-(4-methylbenzoyl)-5H-cyclohepta[b]pyridine

1-(6,7,8,9-Tetrahydro-3-methyl-5H-cyclohepta[b]pyrid-2-yl)-1-(4-methylphenyl)methanol (2.0g) (prepared according to Example 1) in toluene (50ml) with manganese dioxide (6.1g) was refluxed and water
5 collected in a Dean and Stark apparatus. The cooled solution was filtered and the solvent evaporated under reduced pressure to give a gum. This was dissolved in diethyl ether and treated with ethereal HCl. A solid precipitated and was collected by filtration and dried
10 to give the title compound as the hydrochloride (1.8g 80%) m.p. 172-4°C.

Analysis:

Found C, 72.1; H, 7.1; N, 4.8

C₁₉H₂₁NO. HCl requires C, 72.25; H, 7.0; N, 4.4%EXAMPLE 31-(6,7,8,9-Tetrahydro-5H-cyclohepta[b]pyrid-2-yl)-1-(4-methylphenyl)methanol

6,7,8,9-Tetrahydro-5H-cyclohepta[b]pyridine (7.4g
15 0.05m) in tetrahydrofuran (100ml) was cooled to -10°C and treated with a 1.57M solution of n-butyl lithium in n-hexane (32ml), then stirred at -10°C for a further 0.5 hours. 4-Tolualdehyde (10g 0.083m) in tetrahydrofuran (10ml) was added rapidly and the mixture allowed
20 to warm to room temperature, then water and diethyl ether were added. The combined ether extracts were treated with 2 N hydrochloric acid and the aqueous acid extract which separated was basified with solid sodium

-21-

carbonate and then extracted with chloroform. The combined chloroform solutions were washed with water, dried (Mg SO_4) and evaporated to give an oil. This was purified by chromatography using silica columns eluted with diisopropyl ether or chloroform. The purification was monitored by tlc using diisopropyl ether and chloroform on silica plates or samples were silylated with 1-(trimethylsilyl)imidazole and injected on a g.l.c. (injection temp 200°C) with a 25 metre BPI column, running a temperature programme from 120° to 300°C at $10^\circ\text{C}/\text{min}$, then isothermal and carrier gas helium at 10 psi. The title compound having a retention time 17.32 minutes was isolated and converted into the hydrochloride salt by dissolving in diethyl ether and treating with ethereal HCl (1.5g mp $168-170^\circ\text{C}$).

Analysis:

Found C, 70.0; H, 7.4; N, 4.55

$\text{C}_{18}\text{H}_{21}\text{NO} \cdot \text{HCl}$, 0.25 H_2O requires C, 70.1; H, 7.4; N, 4.5%.

EXAMPLE 4

1-(6,7,8,9-Tetrahydro-3-methyl-5H-cyclohepta[b]pyrid-2-yl)-1-phenylmethanol

6,7,8,9-Tetrahydro-3-methyl-5H-cyclohepta[b]pyridine (4.42g, 0.025m) in tetrahydrofuran (50ml) was cooled to -40°C under nitrogen and treated with n-butyl lithium in n-hexane (1.57 molar soln, 16ml) and allowed to warm to 0°C . The solution was cooled to -70°C and treated with a solution of benzaldehyde (4ml) in tetrahydrofuran (10ml) and allowed to warm to room temperature, then water and diethyl ether were added. The organic phase was separated and extracted with 2N hydrochloric acid. The acid phase was made basic with

-22-

solid sodium carbonate and then extracted with chloroform. The combined chloroform extract was washed with water, dried (MgSO_4) and evaporated under reduced pressure to give a gum. This was dissolved in n-hexane
5 and the crystallised material collected by filtration. This was purified by chromatography on silica using diisopropyl ether: chloroform (9:1 by volume) as eluent. The isolated solid was triturated in diisopropyl ether give 1-(6,7,8,9-tetrahydro-3-methyl-
10 5H-cyclohepta[b]pyrid-2-yl)-1-phenylmethanol (1.1g) m.p. 130-2°C.

Analysis:

Found C, 80.95; H, 8.2; N, 5.2

$\text{C}_{18}\text{H}_{21}\text{N}$ requires C, 80.9; H, 7.9; N, 5.2%.

EXAMPLE 5

6,7,8,9-Tetrahydro-3-methyl-2-(1-methyloxy-4-methylbenzyl)-5H-cyclohepta[b]pyridine

1-(6,7,8,9-Tetrahydro-3-methyl-5H-cyclohepta[b]pyrid-
15 2-yl)-1-(4-methylphenyl)methanol (2.0g 0.007m) was added to sodium hydride (0.35g, 50% dispersion in oil, 0.007M, washed with 40-60 petrol) suspended in dimethylformamide (20ml). After evolution of hydrogen had ceased, methyl iodide (0.5ml) was added and the
20 mixture stirred at room temperature for 0.5 hours. Water was added and the mixture extracted with diethyl ether. The combined ether extracts were treated with 2N hydrochloric acid. The aqueous acid solution was separated, basified with solid sodium carbonate and
25 extracted with chloroform. The combined chloroform extracts were washed with water, dried (MgSO_4) and evaporated. The residue was purified by chromatography on silica and eluted with chloroform to give the title compound as a gum. This was dissolved in diethyl

-23-

ether and treated with ethereal HCl to give the crystalline hydrochloride salt, 1.4g m.p. 144-6°C.

Analysis:

Found C, 72.1; H, 7.9; N, 4.4

C₂₀H₂₅ NO. HCl requires C, 72.4; H, 7.9; N, 4.2%.

EXAMPLE 6

1-(6,7,8,9-Tetrahydro-3-methyl-5H-cyclohepta[b]pyrid-2-yl)-1-methyl-1-(4-methylphenyl)methanol

- 5 To a 1.56M solution of n-butyl lithium in n-hexane (13ml) in toluene (20ml) at -20°C under nitrogen, was added a solution of 2-bromo-3-methyl-6,7,8,9-tetrahydro-5H-cyclohepta[b] pyridine (4.8g 0.02m) in toluene (20ml) and kept at -20°C for 0.5 hours. A
10 solution of 4'-methylacetophenone (3.0g 0.022m) was added and the mixture allowed to warm to room temperature and water added. The organic phase was separated and extracted with 2 N hydrochloric acid. This was basified with solid sodium carbonate, then extracted
15 with chloroform. The combined chloroform extracts were dried(MgSO₄) and evaporated under reduced pressure. The residue was purified by chromatography on silica using chloroform as eluent to give a gum. This was dissolved in ether and treated with ethereal HCl to
20 give the title compound as the hydrochloride, three quarter hydrate salt (1.95g) m.p. 162-4°C.

Analysis:

Found C, 69.2; H, 8.0; N, 4.1

C₂₀H₂₅NO. HCl. 0.75H₂O requires C, 69.55; H, 7.7; N, 4.1%.

-24-

EXAMPLE 76,7,8,9-Tetrahydro-3-methyl-2-(4-methylphenyl)-5H-cyclohepta[b]pyridine

1-(6,7,8,9-Tetrahydro-3-methyl-5H-cyclohepta[b]pyrid-2-yl)-1-(4-methylphenyl)methanol (3.0g) was dissolved in ethanol (150ml) and added to 10% Pd/C (1g) under nitrogen. 2 M Sulphuric acid (5ml) was added and the mixture hydrogenated at 50 psi until hydrogen uptake
5 had ceased. The catalyst was removed by filtration and the ethanol removed under reduced pressure. The residue was treated with sodium carbonate solution and then extracted with chloroform. The combined
10 chloroform extracts were dried (MgSO_4) and evaporated to give an oil. The product was separated from starting material by purification using an alumina column eluted with chloroform. The resulting oil was dissolved in diethyl ether and acidified with ethereal
15 HCl to give the title compound as the hydrochloride, (0.47g,) m.p 172-4°C.

Analysis:

Found C, 75.3; H, 7.8; N, 4.4.

C₁₉ H₂₃ N.HCl requires C, 75.6; H, 8.0; N, 4.6%.EXAMPLE 82-(6,7,8,9-Tetrahydro-5H-cyclohepta[b]pyrid-2-yl)-1-(4-methylphenyl)ethanol

To a solution of 1.6M n-butyl lithium in n-hexane (7.2ml) in toluene (25ml), at -20°C under nitrogen, was
20 added a solution of 2-methyl-6,7,8,9-tetrahydro-5H-cyclohepta[b]pyridine (1.7g 0.01m) in toluene (5ml). The solution was left to stir at -20°C for 0.75 hours, then blown over into a solution of 4-methylbenzaldehyde

-25-

(1.5g). The solution was allowed to warm to room temperature and water added. The toluene phase was extracted with 2N hydrochloric acid and this was basified with solid potassium carbonate and then
5 extracted with dichloromethane. The combined dichloromethane extracts were washed with water, dried (MgSO_4) and evaporated to give an oil. This was purified by chromatography on basic alumina using diisopropyl ether as eluent to give the title compound
10 as an oil. This was dissolved in ether and treated with ethereal HCl to give a crystalline hydrochloride salt (1.2g) m.p. 184-6°C.

Analysis:

Found C, 71.7; H, 7.6; N, 4.2

$\text{C}_{19}\text{H}_{23}\text{NO}$, HCl requires C, 71.8; H, 7.6; N, 4.4%.

EXAMPLE 9

1-(6,7,8,9-Tetrahydro-3-methyl-5H-cyclohepta[b]pyrid-2-yl)-1-(4-methylphenyl)methylamine

- 15 a) To a solution of 1.6M n-butyl lithium in n-hexane (16.5ml) in toluene (20ml) at -20°C under nitrogen was added a solution of 2-bromo-6,7,8,9-tetrahydro-3-methyl-5H-cyclohepta[b]pyridine (6.0g, 0.02m) in toluene (30ml). The reaction mixture was kept at -20°C
20 for 0.25 hours and then blown over into a solution of 4-methyl benzonitrile (3.2g) in toluene (30ml) at -20°C. Then mixture was allowed to warm to room temperature. The solvent was removed under reduced pressure to give 1-(6,7,8,9-tetrahydro-3-methyl-
25 5H-cyclohepta[b]pyrid-2-yl)-1-(4-methylphenyl)-methylamine.
- b) Ethanol (150ml) was added to the product of step (a) followed by small portions of sodium borohydride

-26-

(1.2g). After stirring at room temperature for 0.75 hours, 2 N hydrochloric acid was added until hydrogen evolution had ceased. The solvent was removed under reduced pressure and the residue treated with sodium bicarbonate solution and extracted with chloroform. The combined chloroform extracts were washed with water, dried (MgSO_4) and evaporated. The residue was purified by chromatography on basic alumina, eluted with chloroform to give an oil. This was dissolved in diethyl ether and treated with ethereal HCl to give the title compound as the dihydrochloride, monohydrate salt, (1.93g)m.p. 178-180°C.

Analysis:

Found C, 61.7; H, 7.3; N, 7.8

$\text{C}_{19}\text{H}_{24}\text{N}_2 \cdot 2\text{HCl} \cdot \text{H}_2\text{O}$ requires C, 61.45; H, 7.6; N, 7.5%.

EXAMPLE 10

1-(3-Chlorophenyl)-1-(6,7,8,9-tetrahydro-3-methyl-5H-cyclohepta[b]pyrid-2-yl)methanol

To a solution of 1.6M n-butyl lithium in n-hexane (16.5ml) in toluene (20ml) at -20°C under nitrogen was added a solution of 2-bromo-6,7,8,9-tetrahydro-3-methyl-5H-cyclohepta[b]pyridine (6.0g, 0.025m) in toluene (30ml). The reaction mixture was kept at -20°C for 0.25 hours and then blown over into a cooled solution of 3-chlorobenzaldehyde (3.7g) in toluene (30ml) at -20°C, and allowed to warm to room temperature. Water was added and the separated organic phase extracted with 2N hydrochloric acid. This was basified with solid potassium carbonate and extracted with chloroform. The combined chloroform extracts were washed with water, dried (MgSO_4) and evaporated. The solid was purified by passing through a short silica column, eluted with chloroform, then re-crystallised

-27-

from diethyl ether to give the title compound
(4.31g) m.p. 123-5°C.

Analysis:

Found C, 71.4; H, 6.8; N, 5.0

C₁₈H₂₀ClNO requires C, 71.6; H, 6.7; N, 4.6%.

EXAMPLE 11

1-(6,7,8,9-Tetrahydro-3-methyl-5H-cyclohepta[b]pyrid-2-yl)-1-(3-methoxyphenyl)methanol

- 5 To a solution of 1.6M n-butyl lithium in n-hexane
(16.5m,) in toluene (20ml) at -20°, under nitrogen, was
added a solution of 2-bromo-6,7,8,9-tetrahydro-3-methyl
5H-cyclohepta[b]pyridine (6g, 0.025m) in toluene
(30ml). The reaction mixture was kept at -20°C for
10 0.25 hours and then blown over into a cooled solution
(-20°C) of 3-methoxybenzaldehyde (3.7g) in toluene
(30ml) and allowed to warm up to room temperature.
Water was added and the organic phase separated and
treated with 2N hydrochloric acid. A precipitate
15 formed and this was collected by filtration and treated
with sodium carbonate solution, then chloroform. The
organic phase was washed with water, dried (MgO₄) and
evaporated. Trituration with diethyl ether gave
1-(6,7,8,9-tetrahydro-3-methyl-5H-cyclohepta[b]pyrid-2-
20 yl)-1-(3-methoxyphenyl)methanol (4.42g, 59%) m.p.
104-6°C

Analysis:

Found C, 76.9; H, 7.8; N, 4.85

C₁₉H₂₃NO₂ requires C, 76.7; H, 7.8; N, 4.7%.

EXAMPLE 121-(4-Trifluoromethylphenyl)-1-(6,7,8,9-tetrahydro-3-methyl-5H-cyclohepta[b]pyrid-2-yl)methanol

To a solution of 1.6M n-butyl lithium in n-hexane (16.5ml) in toluene (20ml) at -20°C, under nitrogen, was added a solution of 2-bromo-6,7,8,9-tetrahydro-3-methyl-5H-cyclohepta[b]pyridine (6.0g, 0.025m) in
 5 toluene (30ml). The reaction mixture was kept at -20°C for 0.25 hours and then blown over into a solution of a,a,a-tri-fluoro-p-tolualdehyde (4.7g) in toluene (30ml) at -20°C. The solution was allowed to warm up to room temperature and water added. The organic phase
 10 was separated with 2N hydrochloric acid. This was basified with solid potassium carbonate and extracted with chloroform. The combined chloroform extracts were dried (MgSO₄) and evaporated. The residue was recrystallised from n-hexane to give the title
 15 compound, (5.05g) m.p. 31-3°C.

Analysis:

Found C, 68.0; H, 5.9; N, 4.5

C₁₉H₂₀F₃NO requires C, 68.05; H, 6.0; N, 4.2%.

EXAMPLE 131-(6,7,8,9-Tetrahydro-3-methyl-5H-cyclohepta[b]pyrid-2-yl)-1-(pyrid-3-yl)methanol

To a solution of 1.6M n-butyl lithium in n-hexane (16.5ml) in toluene (20ml) at -20°C, under nitrogen,
 20 was added a solution of 2-bromo-6,7,8,9-tetrahydro-3-methyl-5H-cyclohepta[b]pyridine (6g, 0.025m) in toluene (30ml) and kept at -20°C for 0.25 hours. This was blown over into a solution of 3-pyridine carboxaldehyde (2.7g) in toluene (30 ml) kept at -20°C.

The solution was allowed to warm up to room temperature and water added. The organic phase was separated, dried (MgSO_4) and evaporated. The solid was recrystallised from ether (twice) to give 1-(6,7,8,9-tetrahydro-3-methyl-5H-cyclohepta[b]pyrid-2-yl)-1-(pyrid-3-yl)methanol (4.32g, 64%) m.p. 124-6°C.

Analysis:

Found C, 75.8; H, 7.5; N, 10.4.

$\text{C}_{17}\text{H}_{20}\text{N}_2\text{O}$ requires C, 76.1; H, 7.5; N, 10.4%.

EXAMPLE 14

1-(6,7,8,9-Tetrahydro-3-methyl-5H-cyclohepta[b]pyrid-2-yl)-1-(thiophen-2-yl)methanol

To a solution of 1.6M n-butyl lithium in n-hexane (16.5ml) in toluene (20ml) at -70°C, under nitrogen, was added a solution of 2-bromo-6,7,8,9-tetrahydro-3-methyl-5H-cyclohepta[b]pyridine (6g 0.025m) and the mixture kept at -20°C for 0.25 hours. This was blown over into a solution of 2-thiophenecarboxaldehyde (3.0g) in toluene (30ml) kept at -20°C, then allowed to warm to room temperature and water added. The organic phase was extracted with 2N hydrochloric acid, then basified with solid potassium carbonate and extracted with chloroform. The combined chloroform extracts were dried (MgSO_4) and evaporated. The residue was recrystallised from n-hexane to give the title compound (4.42g), m.p. 102-4°C.


Analysis:

Found C, 70.4; H, 7.15; N, 5.3

$\text{C}_{16}\text{H}_{19}\text{NOS}$ requires C, 70.3; H, 7.0; N, 5.1%.

EXAMPLES 15 -20

Using a procedure analogous to Example 6 the following compounds of formula I are prepared according to the reaction scheme



<u>Example No</u>	<u>n</u>	<u>R⁸</u>	<u>R¹</u>	<u>R²</u>	<u>mp</u>
15	3	Me	Me	H	123-4°C
5 16	4	Me	Me	H	88-90°C
17	6	Me	Me	H	118-120°C
18	5	Et	Me	H	
19	5	H	H	Me	
20	5	H	H	H	

-31-

The following examples illustrate the new route to starting materials of formula (IV) used in some of the aforementioned Examples:

Example 21

2-Bromo-3-methyl-6,7,8,9-tetrahydro-5H-cyclohepta[b]-pyridine

5 a) Cycloheptanone was converted to an enamine by reaction with pyrrolidine in toluene solvent and in the presence of toluenesulphonic acid. The enamine was alkylated with methacrylonitrile in ethanol solvent and hydrolysed using acetic acid according to the procedure of Stark et al, J. Amer. Chem. Soc., 85, 207 (1963) to
10 give 2-(2-cyanopropyl)cycloheptanone (bp 100°C/0.02mmHg).

b) The product of step (a) (88g, 0.5m) in acetic acid (1l) was treated with bromine (26ml, 0.5m) over 15 minutes, maintaining an internal temperature of ca,
15 15°C. The mixture was allowed to warm to room temperature overnight. The solvent was evaporated off under reduced pressure and the residue partitioned between water and diisopropyl ether. The organic phase was dried (MgSO₄) and evaporated to give a residue
20 which on recrystallising from hexane gave the title compound as white needles, mp 68-70.

-32-

Examples 22-24

By a process analogous to Example 21 using the appropriate cycloalkanone the following compounds of formula IV wherein p is 0 and Z is Br were prepared:

22. 2-Bromo-3-methyl -5,6,7,8-tetrahydroquinoline,
5 m.p. 50-2°C.

23. 2-Bromo-3-methyl-5H-cyclopenta[b]pyridine.


24. 2-Bromo-3-methyl-5H-cycloocta[b]pyridine.

As illustrated in Example 21 compounds of formula VII may be prepared from cyclic ketone precursors via
10 enamines by Michael alkylation to introduce the
-CHR²CHR¹CN moiety.

-33-


CLAIMS

1. A compound of formula



(I)

or a salt thereof, wherein R¹ and R² each independently represent hydrogen, lower alkyl, lower alkoxy, carboxyloweralkyl, carboxy, hydroxyloweralkyl, halogen, haloloweralkyl, lower alkoxy carbonyl, optionally substituted aryl or optionally substituted aralkyl, n represents an integer from 3 to 6; R³ represents hydrogen or single or multiple substitution on one or more of the aliphatic carbons by one or more substituents selected from lower alkyl, optionally substituted aryl and optionally substituted aralkyl; A represents a group of formula (i) or (ii) below:



in which R⁴, R⁶ and R⁷ each independently represent hydrogen or lower alkyl (providing that when R⁵ is NH₂, R⁴ is hydrogen); m is 0 or 1; R⁵ represents

hydrogen, NH_2 , OH or loweralkoxy, and X is =O, =NH or =NOH, the attachment of A to B being from either end, and B represents an optionally substituted aryl or heteroaryl radical.

2. A compound as claimed in Claim 1 wherein B represents phenyl, pyridyl or naphthyl or such a group substituted by one or more substituents selected from lower alkyl, lower alkoxy, halogen, haloloweralkyl, nitro, amino, lower alkylamino, cyano, carboxy, lower alkoxycarbonyl and lower alkylcarbonyl.

3. A compound as claimed in Claim 1 or Claim 2 in which A is $-\text{CHOH}-$, $-\text{CO}-$, $-\text{CH}_2\text{CO}-$, $-\text{COCH}_2-$, $-\text{CH}(\text{OCH}_3)-$, $-\text{CHOHCH}_2-$, $-\text{CH}_2\text{CHOH}-$, $-\text{CH}_2\text{CH}(\text{OCH}_3)-$ or $-\text{CH}(\text{OCH}_3)\text{CH}_2-$.


4. A compound as claimed in any one of Claims 1 to 3 in which n is 5.

5. A compound as claimed in any one of Claims 1 to 4 in which R^1 and R^2 independently represent hydrogen, lower alkyl, carboxy, lower alkoxycarbonyl, hydroxymethyl, bromine, chlorine, trifluoromethyl, methoxy, phenyl or benzyl.

6. A compound as claimed in any one of Claims 1 to 5 wherein R^3 represents one or more of the following hydrogen, methyl, ethyl, methoxy, ethoxy, phenyl, benzyl or phenyl or benzyl each substituted by halogen, lower alkyl or lower alkoxy.

-35-

7. A compound of formula



(1d)

or a salt thereof,

wherein n represents 4 or 5; R^8 represents lower alkyl, halogen, carboxy, loweralkoxy; R^9 represents hydrogen or a lower alkyl group; A^1 represents $-CO-$ or $-CHOH-$ and B^1 represents phenyl or phenyl substituted by one or more substituents selected from lower alkyl, halogen, lower alkoxy and carboxy.

8. 1-(6,7,8,9-Tetrahydro-3-methyl-5H-cyclohepta[b]-pyrid-2-yl)-1-(4-methylphenyl)methanol or a pharmaceutically acceptable salt thereof.

9. 6,7,8,9-Tetrahydro-3-methyl-2-(4-methylbenzoyl)-5H-cyclohepta[b]pyridine or a pharmaceutically acceptable salt thereof.

10. 1-(6,7,8,9-Tetrahydro-5H-cyclohepta[b]pyrid-2-yl)-1-(4-methylphenyl)methanol or a pharmaceutically acceptable salt thereof.

11. 1-(6,7,8,9-Tetrahydro-3-methyl-5H-cyclohepta[b]-pyrid-2-yl)-1-phenylmethanol or a pharmaceutically acceptable salt thereof.

12. 6,7,8,9-Tetrahydro-3-methyl-2-(1-methoxy-4-methylbenzyl)-5H-cyclohepta[b]pyridine or a pharmaceutically acceptable salt thereof.

13. 2-(6,7,8,9-Tetrahydro-5H-cyclohepta[b]pyrid-2-yl)-1-(4-methylphenyl)ethanol or a pharmaceutically acceptable salt thereof.

14. 1-(3-chlorophenyl)-1-(6,7,8,9-tetrahydro-3-methyl-5H-cyclohepta[b]pyrid-2-yl)methanol or a pharmaceutically acceptable salt thereof,

15. 1-(6,7,8,9-Tetrahydro-3-methyl-5H-cyclohepta[b]pyrid-2-yl)-1-(3-methoxyphenyl)methanol or a pharmaceutically acceptable salt thereof.

16. 1-(4-Trifluoromethylphenyl)-1-(6,7,8,9-tetrahydro-3-methyl-5H-cyclohepta[b]pyrid-2-yl)methanol or a pharmaceutically acceptable salt thereof.

17. 1-(4-Methylphenyl)-1-(6,7-dihydro-3-methyl-5H-cyclopenta[b]pyrid-2-yl)carbinol or a pharmaceutically acceptable salt thereof.

18. 1-(6,7,8,9-Tetrahydro-3-methyl-5H-cyclohepta[b]pyrid-2-yl)-1-methyl-1-(4-methylphenyl)methanol or a pharmaceutically salt thereof.

19. 6,7,8,9-Tetrahydro-3-methyl-2-(4-methylphenyl)-5H-cyclohepta[b]pyridine or a pharmaceutically salt thereof.

20. 1-(6,7,8,9-Tetrahydro-3-methyl-5H-cyclohepta[b]-

-37-

pyrid-2-yl)-1-(4-methylphenyl)methylamine or a pharmaceutically acceptable salt thereof.

21. 1-(6,7,8,9-Tetrahydro-3-methyl-5H-cyclohepta[b]pyrid-2-yl)-1-(pyrid-3-yl)methanol or a pharmaceutically salt thereof.

22. 1-(6,7,8,9-Tetrahydro-3-methyl-5H-cyclohepta[b]pyrid-2-yl)-1-(thien-2-yl)methanol or a pharmaceutically salt thereof.


23. 1-(4-Methylphenyl)-1-(5,6,7,8-tetrahydro-3-methylquinolin-2-yl)carbinol or a pharmaceutically acceptable salt thereof.

24. 1-(4-Methylphenyl)-1-(5,6,7,8,9,10-hexahydro-3-methyl-5H-cycloocta[b]pyrid-2-yl)carbinol or a pharmaceutically salt thereof.

25. A compound as claimed in any one of Claims 1 to 24 when in the form of a salt with an acid selected from hydrochloric, hydrobromic, hydroiodic, sulphuric, phosphoric, nitric, acetic, citric, tartaric, fumaric, succinic, malonic, formic, maleic, methanesulphonic and p-toluenesulphonic acid.


26. A process for preparing a compound of formula I as defined in Claim 1 which comprises one of the following:

a) reacting a compound of formula



(II)

optionally protected in the epi position, wherein R^1 , R^2 , R^3 and n are as defined in Claim 1, p is 0 or 1 and M is sodium, potassium, lithium, M_{hal} where hal is halogen providing that hal is not chlorine when p is 0, with a compound of formula




(III)

wherein B , R^3 , R^6 and R^7 are as defined above and q is 0 or 1 providing that $p + q$ is 0 or 1, followed by protonation and removal of any protecting group present to give a compound of formula I wherein


A is $-CR^4R^5-(CR^6R^7)_m$ where R^5 is OH ;

or b) reacting a compound of formula:



(V)

(if required protected in the epi position) wherein n, R¹, R², R³, R⁶ and R⁷ are as defined in Claim 1, p is 0 or 1 and M¹ is lithium, sodium, potassium or Mghal where hal is chlorine, bromine or iodine, with a nitrile of formula:



(VI)

wherein B, R⁶ and R⁷ are as defined above and q is 0 or 1 providing that p + q = 0 or 1 and if required removing any protecting group to give a compound of formula I wherein A is -CX-(CR⁶R⁷)_m and X is =NH; or

(c) reducing a compound of formula I wherein X is =NH to give a corresponding compound of formula I wherein R⁵ is amino; or

(d) hydrolysing a compound of formula I wherein X is =NH or =NOH to give a compound of formula I wherein X is =O

or

(e) oxidising a compound of formula I wherein R^4 is hydrogen and R^5 is OH to give a compound of formula I wherein X is=O;

or

(f) reacting a compound of formula I wherein X is =O, with hydroxylamine to give a compound of formula I wherein X is =NOH;

or

(g) etherifying a compound of formula I wherein R^5 is OH to give a compound of formula I wherein R^5 is lower alkoxy;

or

(h) hydrogenating a compound of formula I wherein R^5 is OH to give a compound of formula I wherein R^5 is hydrogen,

or


(i) converting a compound of formula I to an acid addition salt.

27. A process for preparing a compound of formula I as claimed in Claim 26 substantially as hereinbefore described and illustrated in any one of Examples 1(b), 2, 3, 4, 5, 6, 7, 8, 9(a), 9(b), 10, 11, 12, 13, 14, 15, 16 and 17.

28. A compound of formula I whenever prepared by a process as claimed in Claim 26 or Claim 27.

29. A pharmaceutical composition comprising a compound of formula I as defined in any one of Claims 1 to 25 or a pharmaceutically salt thereof and a pharmaceutically acceptable carrier.

30. A compound of formula




(IIa)

n, R¹, R², R³, R⁶ and R⁷ are as defined in Claim 1, p is 0 or 1, M is Na, K, Li or MgHal where Hal is a halogen, and P¹ and P² each represent an optional organic silyl protecting group on the carbon epi to the nitrogen.

31. A compound of formula IIa as claimed in claim 30 wherein P¹ and/or P² are/is trimethylsilyl.

32. A process for preparing a compound of formula IIa as claimed in Claim 30 which comprises performing a metallation reaction on a corresponding compound of formula IV




(IV)

if required protected in the epi position by P¹ and/or


P^2 , where Z is hydrogen or halogen and n, p, R^1 , R^2 , R^3 , R^6 and R^7 are as defined in Claim 30, using an appropriate reagent to introduce M.

33. A process for preparing a compound of formula IV



(IV)

where p is 0 and Z is bromine which comprises reacting a compound of formula VII:



(VII)

wherein n, R^1 , R^2 and R^3 are as defined in Claim 1 with either HBr and bromine or with bromine in an inert solvent.

34. A process as claimed in Claim 33 in which the inert solvent is an alkanolic acid or halocarbon.

35. A process as claimed in Claim 33 or Claim 34 wherein n is 5.

-43-

36. A process as claimed in Claim 33 substantially as hereinbefore described and illustrated in any one of Examples 21(b), 22, 23 and 24.

Patents Act 1977
Examiner's report to the Comptroller under
Section 17 (The Search Report)

Application No. 9124015.0
9124015.0

Relevant Technical fields

(i) UK CI (Edition K) C2C CBE CWE CZH

(ii) Int CI (Edition 5) C07D 215/18, 221/04

Databases (see over)

(i) UK Patent Office

(ii) ONLINE DATABASES: CAS ONLINE

Search Examiner

P N DAVEY

Date of Search

5 DECEMBER 1991

Documents considered relevant following a search in respect of claims 1-7

Category (see over)	Identity of document and relevant passages	Relevant to claim(s)
&	GB 2219798 A (WYETH) see whole document	1-7

Category	Identity of document and relevant passages	Relevant to claim(s)

Categories of documents

X: Document indicating lack of novelty or of inventive step.

Y: Document indicating lack of inventive step if combined with one or more other documents of the same category.

A: Document indicating technological background and/or state of the art.

P: Document published on or after the declared priority date but before the filing date of the present application.

E: Patent document published on or after, but with priority date earlier than, the filing date of the present application.

&: Member of the same patent family, corresponding document.

Databases: The UK Patent Office database comprises classified collections of GB, EP, WO and US patent specifications as outlined periodically in the Official Journal (Patents). The on-line databases considered for search are also listed periodically in the Official Journal (Patents).